

Extrasolar Planets

Introduction

So far: have looked at planets around our Sun

Physics question:

Is our Solar System normal?

⇒ Are there planets around other stars?

can then compare solar system with other systems.

To answer these questions, need to detect **extrasolar planets**.

Detection Methods

Possible ways to detect extrasolar planets:

Direct Method:

- ... direct imaging of planet

Indirect Methods: search for evidence for...

- ... gravitational interaction with star in radial velocity
- ... gravitational interaction with star in motion of star
- ... influence of planet on light from behind planet (gravitational lensing)

For time reasons: will look at direct imaging and radial velocity measurements only...

Direct Imaging

In order to make an image of an extrasolar planet, need to **separate images of star and planet** with telescope

⇒ Requires two ingredients:


1. “**contrast**” (relative intensity of star and planet)
2. “**resolving power**” of telescope (angular distance between star and planet)

Direct Imaging: Contrast, I

Estimate intensity contrast between star and planet:

Solar system: **Luminosity** of Sun $L = 3.90 \times 10^{26} \text{ W} =: L_{\odot}$


This power is emitted isotropically into all directions.



Direct Imaging: Contrast, II

Estimate intensity contrast between star and planet:

Solar system: **Luminosity** of Sun $L = 3.90 \times 10^{26} \text{ W} =: L_{\odot}$




This power is emitted isotropically into all directions.

\Rightarrow Energy received per second on whole area of sphere of radius r (area $A = 4\pi r^2$) equals L as well!

Direct Imaging: Contrast, III

Estimate intensity contrast between star and planet:

Solar system: **Luminosity** of Sun $L = 3.90 \times 10^{26} \text{ W} =: L_{\odot}$



This power is emitted isotropically into all directions.

\Rightarrow Energy received per second on whole area of sphere of radius r (area $A = 4\pi r^2$) equals L as well!

\Rightarrow Energy falling per second on area of 1 m^2 at distance r (“**flux**”):

$$F = \frac{L}{4\pi r^2}$$

units: W m^{-2}

Direct Imaging: Contrast, IV

Plugging in typical numbers:

Earth:

distance: $r = 1 \text{ AU} = 150 \times 10^6 \text{ km}$

$\Rightarrow P \sim 1380 \text{ W m}^{-2}$ (“solar constant”).

Total power received by Earth: projected solar facing area

$$A = \pi r_{\oplus}^2 = 1.26 \times 10^{14} \text{ m}^2$$

\Rightarrow Total power received: $P_{\text{total},\oplus} = 1.74 \times 10^{17} \text{ W}$.

Of this, about 30% is reflected, i.e., $L_{\oplus} = 5.2 \times 10^{16} \text{ W} \sim 10^{-10} L_{\odot}$.

The luminosity of the Earth is 10 billion times weaker than that of the Sun.

in infrared, luminosity contrast is only 10 million, but still rather weak. . .

Direct Imaging: Contrast, V

Plugging in typical numbers:

Jupiter:

distance: $r = 5.2 \text{ AU} = 7.8 \times 10^8 \text{ km} \Rightarrow P \sim 51 \text{ W m}^{-2}$

Total power received by Jupiter: projected solar facing area

$$A = \pi r_J^2 = 1.6 \times 10^{16} \text{ m}^2$$

\Rightarrow Total power received: $P_{\text{total}, J} = 8.2 \times 10^{17} \text{ W}$.


Of this, about 30% is reflected, i.e., $L_J = 2.5 \times 10^{17} \text{ W} \sim 6 \times 10^{-10} L_{\odot}$.

The luminosity of Jupiter is ~ 1 billion times weaker than that of the Sun.

\Rightarrow For typical planets in the solar system, **need to be able to detect intensity contrasts of better than 1:1 billion.**

\Rightarrow Not doable now, but not unrealistic to achieve in your lifetime (“coronagraphs”). . .

Direct Imaging: Contrast, VI



How close on sky are images of Sun and planet?

$$\tan \theta = \frac{r}{d} \implies \theta \sim \frac{r}{d}$$

(for small θ : Taylor series: $\tan \theta \sim \theta + (1/3)\theta^3 + \dots$; “small angle approximation”)

Typical distances to nearby stars: $d \sim 100 \text{ Ly} = 9.5 \times 10^{17} \text{ m}$,
 typical distances in planetary system: $r \sim 1 \text{ AU} = 1.5 \times 10^{11} \text{ m}$,

$$\implies \theta = \frac{r}{d} = 1.57 \times 10^{-7} \text{ rad} = 9 \times 10^{-6} \text{ deg} = 0.03''$$

($1'' = 1 \text{ arcsec} = 1/3600 \text{ deg}$).

Direct Imaging: Angular Separation, I

Optics: **resolving power** of telescope with diameter D :

$$\alpha = \frac{12''}{D/1 \text{ cm}}$$

\Rightarrow to resolve $0.03''$, need $D = 4 \text{ m}$

\Rightarrow In principle doable...

Direct Imaging: Angular Separation, II

Optics: **resolving power** of telescope with diameter D :

$$\alpha = \frac{12''}{D/1 \text{ cm}}$$

\Rightarrow to resolve $0.03''$, need $D = 4 \text{ m}$

\Rightarrow In principle doable...

BUT

Earth atmosphere limits resolution to $\sim 0.5''$ (“**seeing**”)

Direct Imaging: Angular Separation, III

Optics: **resolving power** of telescope with diameter D :

$$\alpha = \frac{12''}{D/1 \text{ cm}}$$

⇒ to resolve $0.03''$, need $D = 4 \text{ m}$

⇒ In principle doable...

BUT

Earth atmosphere limits resolution to $\sim 0.5''$ (“**seeing**”)

Currently, direct detection of extrasolar planets not doable, although technologically feasible.

⇒ Need to go to space...

Direct Imaging: Angular Separation, IV

Optics: **resolving power** of telescope with diameter D :

$$\alpha = \frac{12''}{D/1 \text{ cm}}$$

⇒ to resolve $0.03''$, need $D = 4 \text{ m}$

⇒ In principle doable...

BUT


Earth atmosphere limits resolution to $\sim 0.5''$ (“**seeing**”)

Currently, direct detection of extrasolar planets not doable, although technologically feasible.

⇒ Need to go to space...

NASA: **Terrestrial Planet Finder**: 2 missions: 4–6 m telescope (TPF-C; 2014); multiple 3–4 m telescopes (TPF-I, w/ESA; 2020)

ESA: **Darwin**: 6 $> 1.5 \text{ m}$ telescopes, launch planned for 2014



AB Aurigae Disk


Hubble Space Telescope • STIS

PRC99-21 • STScI OPO • C. Grady (NOAO at NASA Goddard Space Flight Center) and NASA

... while planets have yet to be found, direct imaging of the region close to a star is in principle doable with Hubble Space Telescope, but angular resolution not yet good enough.

Radial Velocity Measurements

If we cannot see planet directly \Rightarrow use **indirect methods**.



Two-body problem: **Star** and **planet** move around common **center of mass**:

$$m_1 r_1 = m_2 r_2$$

For circular orbits and orbital period P , velocity of star due to action of planet is

$$v_1 = \frac{2\pi r_1}{P} = \frac{2\pi}{P} \cdot \frac{m_2}{m_1} \cdot r_2$$

Example: Sun vs. Jupiter:

$$m_1 = 2 \times 10^{30} \text{ kg}, m_2 = 2 \times 10^{27} \text{ kg}, r_2 = 5.2 \text{ AU} = 7.8 \times 10^{11} \text{ m}, P_J = 11.9 \text{ yr} = 3.76 \times 10^8 \text{ s}$$


$$\Rightarrow v_1 = 13.1 \text{ m s}^{-1} \sim 50 \text{ km h}^{-1} \sim 30 \text{ mph}$$

Example: Sun vs. Earth gives $v_1 = 10 \text{ cm s}^{-1} \sim 0.8 \text{ km h}^{-1}$

Radial Velocity Measurements

To detect planets, need to be able to measure star velocities with precision to much better than 13 m s^{-1} .


Measure motion of stars using [spectroscopic methods](#).



N.A. Sharp, NOAO/NSO/Kitt Peak FTS/AURA/NSF

Absorption line spectrum of the Sun: Fraunhofer Lines

Radial Velocity Measurements



Using modern spectrographs, position of absorption lines can be measured with very high precision.

Example: H α line from hydrogen in solar spectrum.

but: Light, such as all waves, suffers from **Doppler-effect**: **Lines emitted from moving star are Doppler shifted:**


$$\frac{\lambda_{\text{observed}} - \lambda_{\text{emitted}}}{\lambda_{\text{emitted}}} = \frac{v}{c}$$

\Rightarrow **Can use line shifts to detect extrasolar planets!**

... but need good spectrograph: $v = 13 \text{ m s}^{-1} \Rightarrow \Delta\lambda/\lambda = 4 \times 10^{-8}$, which is only doable by using many tricks.

Results, I

Doppler Shift due to Stellar Wobble




G. Marcy

How to hunt extrasolar planets using the **Doppler Detection Method**:

1. get access to lots of telescope time
2. get access to *very good* spectrograph
3. measure for years, to determine changes in velocity of stars due to motion of star around CM

As of 2005 February, 137 extrasolar planets were known, circling 122 stars.

Results, II




Example: Changing radial velocity of HD 130322 results in discovery of Jupiter-mass planet (Udry et al., 2000).


Here: velocity amplitude: 115 m s^{-1} .

Radial velocity = velocity along our line of sight.

Results, III




G. Marcy/UC Lick



Velocity signature and orbits of the three planets around υ Andromedae.

Results: Mass, I




Many (most!) Planets
found have $M > M_{2+}$

$$M_{2+} = 318 M_{\oplus}$$

Note selection effect: large $M \implies$ larger velocity amplitude
 \implies easier to detect!


So, the fact that we have not seen any Earth-like planets does not mean that they do not exist, just that we cannot detect them yet.



ESA press release, copyright 2002 Lynette Cook, <http://extrasolar.spaceart.org/>.

Jupiter-sized planet plus minor planets (left: detected, Jupiter-scale planet, right: hypothetical planet) around 55 Cancri.


Results: Semimajor Axis, I



Most planets found are close to companion star!


Selection effect: small $a \implies$ short period

\implies detectable in small amount of time (years, not decades)



Jupiter-scale planets close to stars: “hot Jupiters”
e.g., HD 209458b, only
7 Million km from star: planet is
evaporating (HST spectroscopy:
mass loss is 10^7 kg s^{-1})!


Results: Period



Most planets found in short orbits!

Statistics is direct consequence of the selection effect of the previous slide: **short period planets are detectable during typical durations of observing runs...**

Results: Eccentricity



Many planets are in eccentric orbits!


different from solar system!

Might be selection effect due to our existence:

Jupiter in eccentric orbit in our solar system

⇒ strong disturbances of Earth's orbit ⇒ no life!

So, in some sense Copernicus principle does *not* always seem to hold!



D.M.Hardy / PPARC

But not all is bleak – HD 70642 ($d = 90$ ly): discovered by Hugh Jones (Liverpool John Moores University): Jupiter mass planet at 3 AU from solar-like star in circular orbit
 \implies stable Earths are possible.