

Stars

What are stars?

Most important building blocks of the universe: stars

Proper definition:

Stars are gas balls consisting mainly of hydrogen and helium, which produce energy by fusion.

Will now look at observable properties of stars:

1. Distance
2. Luminosity
3. Brightness
4. Masses

... and later deduce how they live from these data.

Distances, I

Direct distance measurements: **parallax measurement**:

⇒ Measure stellar position several times over year with respect to background stars.

Parallax angle (small angle approximation):

$$p = \frac{1 \text{ AU}}{d}$$


(p is measured in *radians*)

Typical values for p are arcseconds

⇒ define distance unit “**Parsec**” (“parallax second”) such that $d = 1 \text{ pc}$ for $p = 1''$:

The parsec is the distance at which 1 AU subtends an angle of $1''$.

Zeilik & Gregory use π instead of p for the parallax...



Distances, II

How far is one parsec?

From $p = 1 \text{ AU}/d$ follows with $p = 1''$:

$$d = 1 \text{ pc} = \frac{1 \text{ AU}}{1''} = \frac{1 \text{ AU}}{\pi/(180 \cdot 3600)} = 206264 \text{ AU} = 3.086 \times 10^{16} \text{ m} \sim 3.26 \text{ ly}$$

Note: If parallax p is known and given in arcseconds, then distance can be immediately computed:

$$\frac{d}{1 \text{ pc}} = \frac{1}{p/1''} \quad \text{or (sloppy notation)} \quad d = \frac{1}{p}$$

Today: positional accuracy $\sim 0.01''$ from ground, and better than 1 mas ($10^{-3}''$) from space \implies can measure parallax out to $\sim 1 \text{ kpc}$

further out: “secondary distance estimators” \implies see later lectures

Luminosity

Definition: **Luminosity** of a star:


The total energy emitted by a star per second is called its **luminosity**.

(=the luminosity is a **power**)

In astronomy, luminosities are often measured in units of the solar luminosity,

$$L_{\odot} = 3.90 \times 10^{26} \text{ J s}^{-1} = 3.90 \times 10^{26} \text{ W}$$

Flux, I



Assumption: star emits its radiation isotropically.

Flux: energy passing per second through area of 1 m^2 at distance r :

$$F = \frac{L}{4\pi r^2}$$

(unit: W m^{-2}).

Flux, II

Fluxes from stars (apart from the Sun) are *very small*.

Example: α Centauri (closest star to the Sun).

- distance: $1.3 \text{ pc} \sim 4 \times 10^{16} \text{ m}$
- luminosity: similar to the Sun ($4 \times 10^{26} \text{ W}$).

Flux, III

Fluxes from stars (apart from the Sun) are *very small*.

Example: α Centauri (closest star to the Sun).

- distance: 1.3 pc $\sim 4 \times 10^{16}$ m

- luminosity: similar to the Sun (4×10^{26} W).

\implies Flux arriving on Earth:

$$F = \frac{L}{4\pi r^2} = \frac{3.9 \times 10^{26} \text{ W}}{4\pi \cdot 16 \cdot 10^{32} \text{ m}^2} = 2 \times 10^{-8} \text{ W m}^{-2}$$

(compare with solar constant, $F = 1380 \text{ W m}^{-2}$!)

Flux, IV

Fluxes from stars (apart from the Sun) are *very small*.

Example: α Centauri (closest star to the Sun).

- distance: 1.3 pc $\sim 4 \times 10^{16}$ m

- luminosity: similar to the Sun (4×10^{26} W).

\implies Flux arriving on Earth:

$$F = \frac{L}{4\pi r^2} = \frac{3.9 \times 10^{26} \text{ W}}{4\pi \cdot 16 \cdot 10^{32} \text{ m}^2} = 2 \times 10^{-8} \text{ W m}^{-2}$$

(compare with solar constant, $F = 1380 \text{ W m}^{-2}$!)

\implies your eye detects a power of

$$P = A_{\text{eye}} F = 5 \times 10^{-12} \text{ W}$$

from α Cen (assuming $A_{\text{eye}} \sim 25 \text{ mm}^2$)!

weakest visible stars: $\sim 100\times$ weaker!

Magnitudes, I



Hipparchus
(??– ~127 BC)

First classification of stars:

- Stars of “magnitude 1”: brightest (visible) stars
- Stars of “magnitude 6”: faintest (visible) stars

<http://www-gap.dcs.st-and.ac.uk/~history/Mathematicians/Hipparchus.html>

Magnitudes, II

Pogson (1865): Eye sensitivity is logarithmic, such that

A brightness *difference* of 5 magnitudes corresponds to a *ratio* of 100 in detected flux

Magnitudes, III

Pogson (1865): Eye sensitivity is logarithmic, such that

A brightness *difference* of 5 magnitudes corresponds to a *ratio* of 100 in detected flux

So, if magnitudes of two stars are m_1 and m_2 , then

$$\frac{f_1}{f_2} = 100^{(m_2 - m_1)/5}$$

Magnitudes, IV

Pogson (1865): Eye sensitivity is logarithmic, such that

A brightness *difference* of 5 magnitudes corresponds to a *ratio* of 100 in detected flux

So, if magnitudes of two stars are m_1 and m_2 , then

$$\frac{f_1}{f_2} = 100^{(m_2 - m_1)/5}$$

This means:

$$\log_{10}(f_1/f_2) = \frac{m_2 - m_1}{5} \log_{10} 100 = \frac{2}{5}(m_2 - m_1)$$

Magnitudes, V

Pogson (1865): Eye sensitivity is logarithmic, such that

A brightness *difference* of 5 magnitudes corresponds to a *ratio* of 100 in detected flux

So, if magnitudes of two stars are m_1 and m_2 , then

$$\frac{f_1}{f_2} = 100^{(m_2 - m_1)/5}$$

This means:

$$\log_{10}(f_1/f_2) = \frac{m_2 - m_1}{5} \log_{10} 100 = \frac{2}{5}(m_2 - m_1)$$

or

$$m_2 - m_1 = 2.5 \log_{10}(f_1/f_2) = -2.5 \log_{10}(f_2/f_1)$$

Note: Larger Magnitude = *FAINTER* Stars

Luminosity (revisited)

Inverse square law links flux f at distance d to flux F measured at another distance D :

$$\frac{F}{f} = \frac{L/4\pi D^2}{L/4\pi d^2} = \left(\frac{d}{D}\right)^2$$

Convention: to describe luminosity of a star, use the **absolute magnitude** M , defined as magnitude measured at distance $D = 10$ pc.

Therefore,


$$m - M = 2.5 \log(F/f) = 2.5 \log(d/10 \text{ pc})^2 = 5 \log d - 5$$


$m - M$ is called the **distance modulus**, d is measured in pc.





V. van Gogh: Starry Night over the Rhône (1888)


The WebMuseum (<http://www.ibiblio.org/wm/>; original: Paris, Musée d'Orsay)


















ζ^1 Ursae Majoris



Masses, X

Mizar A and B are rather typical stars:

50% – 80% of all stars in the solar neighbourhood belong to multiple systems.

Rough classification:

Masses, XI

Mizar A and B are rather typical stars:

50% – 80% of all stars in the solar neighbourhood belong to multiple systems.

Rough classification:

apparent binaries: stars are *not* physically associated, just happen to lie along same line of sight (“**optical doubles**”).

Masses, XII

Mizar A and B are rather typical stars:

50% – 80% of all stars in the solar neighbourhood belong to multiple systems.

Rough classification:

apparent binaries: stars are *not* physically associated, just happen to lie along same line of sight (“**optical doubles**”).

visual binaries: bound system that can be resolved into multiple stars (e.g., Mizar); can **image orbital motion**, **periods typically 1 year to several 1000 years**.

Masses, XIII

Mizar A and B are rather typical stars:





50% – 80% of all stars in the solar neighbourhood belong to multiple systems.

Rough classification:

apparent binaries: stars are *not* physically associated, just happen to lie along same line of sight (“**optical doubles**”).


visual binaries: bound system that can be resolved into multiple stars (e.g., Mizar); can **image orbital motion**, **periods typically 1 year to several 1000 years**.

spectroscopic binaries: bound systems, cannot resolve image into multiple stars, but **see Doppler effect in stellar spectrum**; often **short periods (hours... months)**.



Potpourri of observed binary star orbits.

Masses, XV



Problem when analysing orbits: **orientation of orbit in space**: “**inclination**”

In simplest case: real semimajor axis:

$$a_{\text{observed}} = a_{\text{real}} \cos i$$

Inclination typically found using Kepler’s 2nd law plus geometry...

Masses, XVI

To determine stellar masses, use **Kepler's 3rd law**:

$$\frac{a^3}{P^2} = \frac{G}{4\pi^2}(m_1 + m_2)$$

where

- $M_{1,2}$: masses
- P : period
- a semimajor axis

Masses, XVII

To determine stellar masses, use **Kepler's 3rd law**:

$$\frac{a^3}{P^2} = \frac{G}{4\pi^2}(m_1 + m_2)$$

where

- $M_{1,2}$: masses
- P : period
- a semimajor axis

Observational quantities:

- P – directly measurable
- a – measurable from image *if and only if* distance to binary and the inclination is known

Masses, XVIII

From Kepler's 3rd law, can determine $M_1 + M_2$.


Need to determine individual masses, M_1 and M_2 :

⇒ use center of mass (CM):

$$M_1 a_1 = M_2 a_2 \quad \text{such that} \quad \frac{M_1}{M_2} = \frac{a_2}{a_1}$$

where a_1, a_2 : semi-major axes of orbits around CM (observable from imaging).

Spectroscopic Binaries, I



For **spectroscopic binaries**: can only measure **radial velocity** along line of sight
 For circular orbit, angle θ on orbit:

$$\theta = \omega t$$

where $\omega = 2\pi/P$.

Observed radial velocity:


$$v_r = v \cos(\omega t)$$

If orbit has inclination i , then

$$v_r(t) = v \sin i \cos(\omega t)$$

From observation of $v_r(t) \implies v \sin i$.
 (“**velocity amplitude**”)

Spectroscopic Binaries, II



HDE 226868/Cyg X-1; Pottschmidt (2001)

Motion of star visible through **Doppler shift** in stellar spectrum:

$$\frac{\Delta\lambda}{\lambda} = \frac{v_r}{c} = \frac{v}{c} \sin i \cos \omega t$$

For almost all stars, classical Doppler effect is enough; once $v \gtrsim 0.1c$, however, use **relativistic Doppler effect**,

$$\nu_{\text{obs}} = \nu_{\text{em}} \sqrt{\frac{1 + v/c}{1 - v/c}}$$

Mass Function

If only one star visible: can only determine **limits for mass**: **mass function**

$$\frac{P v_{\text{obs}}^3}{2\pi G} = \frac{M_2^3 \sin^3 i}{(M_1 + M_2)^2} =: f_M$$

(see handout)

with **observables**:

- v_{obs} : (velocity amplitude of M_1)
- P : period

and **unknowns**:

- M_1 : mass of “**primary star**”
- M_2 : mass of (unseen) “**secondary star**”
- i : inclination

$\implies f_M$ is **lower limit for M_2** , since for $M_1 = 0$, $M_2 = f_M / \sin^3 i \geq f_M$

Often used for neutron star and black hole binaries...

Note that the following is for your information only, you will not be tested on your ability to memorize this lengthy derivation. . .

To derive the mass function, we start as usual with Kepler's 3rd law,

$$\frac{G}{4\pi^2}(M_1 + M_2) = \frac{R^3}{P^2}$$

In the following, we will assume that we observe the spectral lines from star number 1 only.

Because of the center of mass definition,

$$M_1 r_1 = M_2 r_2$$

such that

$$R = r_1 + r_2 = r_1 \left(1 + \frac{r_2}{r_1} \right) = r_1 \left(1 + \frac{M_1}{M_2} \right)$$

In the case that the orbits are circular, the velocity of the star whose spectrum we see is

$$v_1 = \frac{2\pi r_1^2}{P}$$

However, due to the unknown inclination, we only observe the radial velocity component, that is

$$v_{\text{obs}} = v_1 \sin i$$

In terms of the observables, r_1 is

$$r_1 = \frac{P}{2\pi} v_1 = \frac{P}{2\pi} \frac{v_{\text{obs}}}{\sin i}$$

such that finally

$$R = r_1 \left(1 + \frac{M_1}{M_2} \right) = \frac{P}{2\pi} \frac{v_{\text{obs}}}{\sin i} \left(1 + \frac{M_1}{M_2} \right)$$

We can now insert R into Kepler's 3rd law:


$$\frac{G}{4\pi^2}(M_1 + M_2) = \frac{1}{P^2} \frac{P^3}{(2\pi)^3} \frac{v_{\text{obs}}^3}{\sin^3 i} \left(1 + \frac{M_1}{M_2} \right)^3$$

and obtain after some straightforward algebra

$$\frac{M_2^3}{(M_1 + M_2)^2} \sin^3 i = \frac{P v_{\text{obs}}^3}{2\pi G}$$

the mass function. On the right side are the observables P and v_{obs} , on the left hand side the unknowns i , M_1 , and M_2 .


Application: Mass-Luminosity Relation



Can now look at solar neighborhood:

- **apparent magnitude** m *and* **distance** \implies **luminosity**
 - **mass** from binary stars
- \implies determine **mass-luminosity relationship**

Application: Mass-Luminosity Relation



sometimes, one also sees $L \propto M^{3.3} \dots$

Empirical result:

$$\frac{L}{L_{\odot}} = \begin{cases} 0.23 \left(\frac{M}{M_{\odot}} \right)^{2.3} & (M < 0.43 M_{\odot}) \\ \left(\frac{M}{M_{\odot}} \right)^{4.0} & (M \geq 0.43 M_{\odot}) \end{cases}$$

\implies more massive stars have extremely higher luminosities!

(factor 2 in $M \rightarrow$ factor 8 in L).

Direct consequence:

More massive stars live much shorter lives

Introduction

Final observable of stars: **Temperature**

Obtained using **spectroscopy**

In the following: rough outline, as stellar spectroscopy is rather complicated
⇒ see module “stars” for the g(l)ory details...

Outline:

1. **Planck’s Radiation Laws**
2. **Stellar Continuum Spectra**
3. **Spectral Classification**

... unfortunately, need to be a little bit formal first

Planck's Radiation Law, I



Max Planck (1858–1947)

Stars are big glowing gas balls.

In zeroth order: thermodynamic equilibrium.


Max Planck: under these circumstances:
emitted spectrum is blackbody radiation:

$$F_{\lambda} = \frac{2hc^2/\lambda^5}{\exp(hc/\lambda kT) - 1}$$

F_{λ} : Energy emitted per second and wavelength interval

- $h = 6.623 \times 10^{-34}$ J s: Planck's constant
- $k = 1.38 \times 10^{-23}$ J K⁻¹: Boltzmann constant

Planck's Radiation Law, II



Without proof, the following two important relationships hold for blackbody radiation:

Stefan-Boltzmann law: Power emitted per square-metre surface of a blackbody:

$$P = \sigma T^4$$

where $\sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$
 “hotter bodies have a much higher luminosity”

Wien's displacement law: Wavelength of maximum blackbody emission:

$$\lambda_{\text{max}} T = 2.898 \times 10^{-3} \text{ m K}$$

“hotter bodies radiate higher energetic radiation”


Spectroscopy, I

Quantum mechanics: **atoms have discrete energy levels**


Energy levels in Hydrogen:

$$E_n = -\frac{2\pi^2\mu e^4}{\hbar^2} \cdot \frac{1}{n^2} \propto -\frac{1}{n^2}$$

($n \in \mathbb{N}$; **Balmer formula**)




Spectroscopy, II



In hydrogen atom: electrons typically found in ground state.


if temperature is higher, can also be in 1st excited state, but the physical principles following remain the same....

Spectroscopy, III




Photon hitting atom has energy $E_{\text{phot}} = h\nu = hc/\lambda$. If $E_{\text{phot}} = E_2 - E_1$, then photon can be absorbed...

Spectroscopy, IV



Photon hitting atom has energy $E_{\text{phot}} = h\nu = hc/\lambda$. If $E_{\text{phot}} = E_2 - E_1$, then **photon can be absorbed**. . . and electron has higher energy (is **excited**).

Spectroscopy, V




After absorption event, absorbing photon has disappeared and hydrogen atom remains in **excited state**.

Spectroscopy, VI


1. Assume stellar surface has **continuum spectrum** (Planck).

Emitter of
continuum radiation






Spectroscopy, VII


1. Assume stellar surface has **continuum spectrum** (Planck).
2. Assume surface is below atmosphere of colder gas.



Spectroscopy, VIII




1. Assume stellar surface has **continuum spectrum** (Planck).
 2. Assume surface is below atmosphere of colder gas.
- ⇒ Atmosphere **absorbs photons at wavelengths characteristic for the elements present in stellar atmosphere.**
- ⇒ **Formation of absorption line spectrum.**



N.A.Sharp, NOAO/NSO/Kitt Peak FTS/AURA/NSF

Absorption line spectrum of the Sun: Fraunhofer Lines



Annie Cannon (around 1890): Stars have different spectra.

NOAO

HD 12993

HD 158659

HD 30584

HD 116608

HD 9547

HD 10032

BD 61 0367

HD 28099

HD 70178

HD 23524

SAO 76803

HD 260655

Yale 1755

HD 94082

SAO 81292

HD 13256

Annie Cannon (around 1890): Stars have different spectra.

NOAO

Stellar Spectra




Annie Jump Cannon (1863–1941)

Biography: <http://www.sdsc.edu/ScienceWomen/cannon.html>


Annie Jump Cannon: There are spectral types.

Henry Draper catalogues (Cannon plus ~10 female “computers”): 225000 spectral classifications.







Annie Cannon: Strength of absorption lines varies with spectral type.



Annie Cannon: Strength of absorption lines varies with spectral type.


Stellar Spectra



Silva & Cornell, 1992, ApJ Suppl. 81, 865

Cecilia Payne-Gaposchkin: **Spectral sequence is temperature sequence.**

Stellar Spectra



Cecilia Payne-Gaposchkin (1900–1979)

Biography: <http://www.harvardsquarelibrary.org/unitarians/payne2.html>

BSc, Cambridge, left UK because of situation of women in astronomy

1st person to obtain PhD in Astronomy at Harvard: “Stellar Atmospheres, A Contribution to the Observational Study of High Temperature in the Reversing Layers of Stars”

Otto Struve: “undoubtedly the most brilliant Ph.D. thesis ever written in astronomy.”

Spectral types are a temperature sequence.

later: 1st female full professor at Harvard

Stellar Spectra

Summary spectral classes as a temperature sequence.

O - B - A - F - G - K - M

30000 K

“early type”


3000 K

“late type”

plus subtypes: B0... B9, A0... A9, etc.

Sun is G2.

Note: “early” and “late” has *nothing* to do with age!



Stellar Spectra

Summary spectral classes as a temperature sequence.

O - B - A - F - G - K - M

30000 K

3000 K

“early type”

“late type”

plus subtypes: B0... B9, A0... A9, etc.


Sun is G2.

Note: “early” and “late” has *nothing* to do with age!

Mnemonics:

(<http://lheawww.gsfc.nasa.gov/users/allen/obafgkmrns.html>)

O Be A Fine Girl Kiss Me



Stellar Spectra

Summary spectral classes as a temperature sequence.

O - B - A - F - G - K - M

30000 K

3000 K

“early type”

“late type”

plus subtypes: B0... B9, A0... A9, etc.


Sun is G2.

Note: “early” and “late” has *nothing* to do with age!

Mnemonics:

(<http://lheawww.gsfc.nasa.gov/users/allen/obafgkmrns.html>)

O Be A Fine Girl
 Guy Kiss Me



Stellar Spectra

Summary spectral classes as a temperature sequence.

O - B - A - F - G - K - M

30000 K

3000 K

“early type”

“late type”

plus subtypes: B0...B9,A0...A9, etc.

Sun is G2.


Note: “early” and “late” has *nothing* to do with age!

Mnemonics:

(<http://lheawww.gsfc.nasa.gov/users/allen/obafgkmrns.html>)

O Be A Fine
Guy Kiss Me

Only Boys Accepting Feminism Get Kissed Meaningfully



Stellar Spectra

Summary spectral classes as a temperature sequence.

O - B - A - F - G - K - M

30000 K

3000 K

“early type”

“late type”

plus subtypes: B0... B9, A0... A9, etc.

Sun is G2.

Note: “early” and “late” has *nothing* to do with age!


Mnemonics:

(<http://lheawww.gsfc.nasa.gov/users/allen/obafgkmrns.html>)


O Be A Fine Girl
 Guy Kiss Me

Only Boys Accepting Feminism Get Kissed Meaningfully

Only Bold Astronomers Forge Great Knowledgeable Minds




HRD



Hertzsprung - Russell Diagram (HRD): Stellar temperature versus stellar luminosity


HRD



- Most stars on **Main Sequence** (“dwarfs”)

Hertzsprung - Russell Diagram (HRD): Stellar temperature versus stellar luminosity

HRD




- Most stars on **Main Sequence** (“dwarfs”)
- Stellar Luminosity:

$$L = 4\pi R^2 \sigma T^4 \propto R^2 T^4$$
 \implies cold, luminous stars are *BIG*
 \implies “giants”

Hertzsprung - Russell Diagram (HRD): Stellar temperature versus stellar luminosity

HRD




- Most stars on **Main Sequence** (“dwarfs”)
- Stellar Luminosity:

$$L = 4\pi R^2 \sigma T^4 \propto R^2 T^4$$
 \implies cold, luminous stars are **BIG**
 \implies “giants”
- Hot, underluminous stars are small: “white dwarfs”

Hertzsprung - Russell Diagram (HRD): Stellar temperature versus stellar luminosity

HRD



Combining Mass-Luminosity Relationship and HRD:


Main Sequence is a Mass Sequence

- M-Dwarfs have $M \lesssim 0.25M_{\odot}$
- G-Stars are similar to Sun and have $M \sim M_{\odot}$
- O- and B-Stars are very massive ($M \gtrsim 20M_{\odot}$)




Globular Cluster NGC 6903

STScI



M3, S. Kafka and K. Honeycutt, Indiana University/WIYN/NOAO/NSF
note many red giants!



HRD of Globular Cluster M5 (UNSW, Sydney)
 (B-V: \sim spectral class; V is a magnitude)

Globular Clusters: HRD is very different of solar neighbourhood

MS: Main Sequence

TO: Turn-Over point

HB: Horizontal Branch

RGB: Red Giant Branch

AGB: Asymptotic Giant Branch

WD: White Dwarfs

All stars in globular cluster born at the same time

\implies HRD shows evidence for stellar evolution


Stellar Evolution



Orion Nebula; R. Gendler



Optical View of B68 (ESO; VLT/FORS1)



IR View of B68 (ESO; VLT/FORS1 + NTT/SOFI)

Stellar Birth, V

Stars are born in “Giant Molecular Clouds”.

Pieces of cloud **collapse**, if mass within radius of cloud is larger than **Jeans mass**:

$$M_J \sim \frac{4\pi}{3} R_J^3 \rho$$

... which has typical values of 50–100 M_\odot .

After this:

⇒ **collapsing piece fragments** into smaller pieces of $\sim 1 M_\odot$ mass (takes $< 10^6$ years)

⇒ **density** in centre of each piece **increases**

⇒ $T \gtrsim 4 \times 10^6$ K

⇒ **nuclear fusion starts**

⇒ **star is formed**

The following is for your information only and will not be assessed in any way:

In order to derive the Jean's mass, let's look at a simple model where a cloud in the interstellar medium is kept together by its own gravitation. Because the cloud has a temperature above absolute zero, thermal motion from the gas particles will try to disperse the cloud. The cloud is thus gravitationally only if the total energy of the gas particles, i.e., the sum of the particle kinetic energy and the gravitational binding energy, is negative:

$$\frac{3}{2} \frac{M}{m_p} kT - \frac{3}{5} \frac{GM^2}{R} \leq 0$$

If the energy is less than zero, the cloud is not in hydrostatic equilibrium and will collapse. This is the case if

$$\frac{M}{R} \geq \frac{5}{2} \frac{kT}{Gm_p} \quad \text{or} \quad \frac{4\pi}{3} \rho R^2 \geq \frac{5}{2} \frac{kT}{Gm_p}$$

This is true for radii greater than the **Jeans length**,

$$R_J = \sqrt{\frac{15kT}{8\pi Gm_p\rho}} \sim \sqrt{\frac{kT}{Gm_p\rho}}$$

Plugging in typical values, i.e., $T \sim 50$ K, particle density $n = 10^5$ H-atoms cm^{-3} (that is a mass density of $\rho = nm_p \sim 1.7 \times 10^{-9}$ g cm^{-3}) gives $R_J \sim 0.2$ pc, corresponding to a Jeans mass of around 70 solar masses.



Tarantula Nebula in Large Magellanic Cloud (MPG/ESO 2.2 m / WFI)



HH34 in Orion (ESO VLT KUEYEN/FORS2)

Main Sequence, I

Once star has collapsed and nuclear fusion has started: **main sequence**


The Main Sequence is the result of steady state fusion (“burning”) of hydrogen into helium in stellar centres.

... longest phase of stellar evolution (10 billion years for Sun)

Stellar structure defined by balance between pressure inwards due to gravitation and pressure outwards due to energy release (“**hydrostatic equilibrium**”).

Main Sequence, II

Nuclear fusion:



How much energy is gained?

Particle physics: express mass as “rest energy equivalent” via $E = mc^2$ (and call it “mass”...).

usually use energy units of MeV, $1 \text{ MeV} = 1.602 \times 10^{-13} \text{ J}$

mass of 4 protons ($4 \times 938 \text{ MeV}$):	3752 MeV
– mass of ${}^4_2\text{He}$:	3727 MeV
mass defect Δmc^2 :	25 MeV

In the fusion of hydrogen to helium, 0.7% of the available rest mass energy is converted to energy.

Nuclear physics: efficiency of H-burning strongly depends on temperature

⇒ explanation for mass-luminosity relation (massive stars have hotter cores)

Main Sequence, III

Stellar structure governed by **four coupled differential equations**:

Mass structure

(mass conservation)

$$\frac{dM}{dr} = 4\pi r^2 \rho(r)$$

Pressure structure

(hydrostatic equilibrium)

$$\frac{dP}{dr} = -\rho(r) \frac{GM(r)}{r^2}$$

Temperature structure

(energy transport)

$$\frac{dT}{dr} = -\frac{3}{4ac} \frac{k\rho(r)}{T^3} \frac{L(r)}{4\pi r^2}$$

Energy conservation

(energy transport)


$$\frac{dL}{dr} = 4\pi r^2 \rho(r) \epsilon(r)$$

plus “equation of state” ($P = P(T, \rho)$), energy generation ($\epsilon = \epsilon(T, \rho, Z)$),...

Stellar model: numerical solution of stellar structure equations.


26 Feb 2004

19:00:29 UT




ISOON - Sunspot

630.32 nm



(c) Scientific American

Main Sequence, VI



after Iben, 1991

Evolution of stars in the HRD from main sequence to death

Typical timescales (units of 10^6 yr; Schaller et al. 1992):

	$1 M_{\odot}$	$5 M_{\odot}$	$25 M_{\odot}$
H \rightarrow He	10000	94	6.4
He \rightarrow C		12	0.6
C+O			0.01
PN	$\lesssim 0.01$	$\lesssim 0.01$	N/A
WD	∞	∞	N/A

Post-H-burning burning: need higher core temperatures (Coulomb barrier!), less energy release \implies last much shorter than hydrogen burning.




Abell 39 (WIYN, AURA, NOAO, NSF)

planetary nebulae: material ejected during AGB phase, photoionized
once central star emits UV photons; after PN phase: white dwarf
End result of stellar evolution for stars with $M \leq 5 M_{\odot}$.



Ring Nebula (HST/STScI/NASA)

planetary nebulae: material ejected during AGB phase, photoionized
once central star emits UV photons; after PN phase: white dwarf
End result of stellar evolution for stars with $M \leq 5 M_{\odot}$.



IC4406 (ESO VLT)

planetary nebulae: material ejected during AGB phase, photoionized
once central star emits UV photons; after PN phase: white dwarf
End result of stellar evolution for stars with $M \leq 5 M_{\odot}$.



NGC 6853/M27 (“Dumbbell Nebula”; ESO VLT/FORS)

planetary nebulae: material ejected during AGB phase, photoionized
once central star emits UV photons; after PN phase: white dwarf
End result of stellar evolution for stars with $M \leq 5 M_{\odot}$.

White Dwarfs

Reminder: stars: hydrostatic equilibrium, **inwards gravitational pressure balanced by outwards gas pressure**

For gas pressure ($P = nkT$): **energy source needed** to heat gas (=fusion).

End of stellar life: energy source ceases to work \implies **gravitational collapse!**

BUT:

collapse cannot continue indefinitely:

increased density \implies **quantum mechanical effects** become important

Different ways to write the equation of state of an ideal gas

Among the more confusing subjects of thermodynamics are the many different ways in which the ideal gas equation can be written.

The one I prefer for astronomy is

$$P = nkT$$

where

- P : Pressure (measured in $N\ m^{-1}$)
- n : particle density (i.e., number of particles per cubic metre, unit: m^{-3})
- $k = 1.38066 \times 10^{-23}\ J\ K^{-1}$: Boltzmann constant
- T : Temperature (measured in Kelvins)

This equation has the advantage that it counts all particles individually (thus using n). If you know the mass of the gas particles, m_{gas} then another way of writing the ideal gas equation is

$$P = \frac{nm_{\text{gas}}}{m_{\text{gas}}}kT = \rho kT \frac{1}{m_{\text{gas}}}$$

illustrating that for an ideal gas, $P \propto \rho$, where ρ is the mass density.

Another way to write the ideal gas equation is in terms of the total number of gas molecules, $N = nV$, where V is the volume. The ideal gas equation then is

$$P = \frac{N}{V}kT \iff PV = NkT$$

This version has the problem, however, that the number of gas molecules is typically rather large (there are 6×10^{23} molecules in a volume of 22.4 litres of gas, this number of particles is called one *mole*). Because working with smaller numbers is generally better, chemists prefer to work with moles. Per definition, the unit of particle number here is the Avogadro number $N_A = 6.0221 \times 10^{23}$. So, if you want to work with moles, then the above equation becomes

$$PV = \frac{N}{N_A}AkT = N_{\text{mol}}RT$$

where

- N_{mol} : the number of moles of the gas in the volume V ,
- $R = N_A k = 8.3145\ J\ mol^{-1}\ K^{-1}$: the universal gas constant

To summarise, each of these equations has its own uses, and which one you want to use, really depends on the circumstances of the problem you are solving. For your future life as physicists, try to remember one of them, and then understand how you get from this one to the others, instead of memorising all four ones. This approach will need less memory and lead to a better understanding of what is really going on behind the scenes.

QM interlude, I

Quantum mechanics: One of the weirder phenomena in QM is the **Pauli exclusion principle**:

For particles such as electrons (“Fermions”), at least one of their quantum numbers must be different.

Quantum numbers are, e.g.,


- position (x, y, z) ,
- momentum (mv_x, mv_y, mv_z) ,
- angular momentum,
- spin (s)

All of these numbers are “**quantized**”, i.e., can only have discrete values (e.g., spin: $+1/2, -1/2$).

In typical gas, this is not a problem (“**phase space is (almost) empty**”), but once it becomes dense \implies exclusion principle kicks in.

QM interlude, II


Effect of high density on electron energy



Energy of electrons at
the same position in
space


QM interlude, III

Effect of high density on electron energy



Energy of electrons at
the same position in
space


QM interlude, IV



Effect of high density on electron energy

Energy of electrons at
the same position in
space


QM interlude, V



Effect of high density on electron energy

Energy of electrons at
the same position in
space


QM interlude, VI



Effect of high density on electron energy

Energy of electrons at
the same position in
space

QM interlude, VII




Energy of electrons at
the same position in
space

Effect of high density on electron energy:

In **degenerate electron gases**, electrons have much higher energies than in thermal gas.

QM interlude, VIII



Energy of electrons at
the same position in
space

Effect of high density on electron energy:


In **degenerate electron gases**, electrons have much higher energies than in thermal gas.

Interaction of electrons results in **degeneracy pressure**:

$$P = \frac{\hbar^2}{m_e} n_e^{5/3} \propto \rho^{5/3}$$

Note: The degeneracy pressure is **independent of the temperature!**

White Dwarfs: Summary




Sirius A+B: *Chandra*
(X-rays; *WD is bright*)



McDonalds Observatory
(optical; *WD is faint*)


White Dwarfs:

1. End stages of evolution of stars with $M \lesssim 10 M_{\odot}$ on main sequence
2. typically $M \sim 0.8 M_{\odot}$, and always $M < 1.44 M_{\odot}$ (**Chandrasekhar mass**; above that: relativistic degenerate gas ($P \propto \rho^{4/3}$), can show that under these circumstances WD is not stable.
3. mainly consist of **C and O**
4. **Radius \sim Earth**
5. Typical **density** $\rho \sim 10^6 \text{ g cm}^{-3}$
6. **interior temperature** $\sim 10^7 \text{ K}$, **atmosphere** $\sim 10^4 \text{ K}$, slowly cooling down (observable for $\gtrsim 10^9$ years).




Type II SN2001cm in NGC5965 (2.56 m NOT, Håkon Dahle; NORDITA)

Evolution of more massive stars: fusion up to ^{56}Fe , then no energy gain
 \implies no pressure balance in centre \implies supernova explosion of type II.
energy release: 10^{46} W ($10^{20} L_{\odot}$; about 1% in light, rest in neutrinos)



(ESO VLT/FORS 2)

Crab nebula: young remnant of SN of 1054, observed light due to **synchrotron radiation** (radiation emitted by electrons accelerated in magnetic field)



5000–10000 year old IC 1340/Veil Nebula/Cygnus Loop (©Loke Kun Tan)

Older supernova remnants: “wispy structure” due to interaction with interstellar medium, radiation (line emission) mainly caused by heating due to shocks.

Neutron Stars

During SN explosion:

Core of exploding star above Chandrasekhar limit \implies core collapses

Densities get so high that neutronization sets in:



General properties:

- Pressure mainly through degenerate neutrons (similar to degenerate electrons for WD!).
- Typical density: $\rho \sim 10^{14} \text{ g cm}^{-3}$ (nuclear densities)
- Typical radius: 10... 15 km (Coventry!)
- surface gravity $\sim 10^{11} \times \text{Earth}$
- Detailed structure not yet fully understood,

Neutron Stars: Rotation, I

During SN collapse, **angular momentum is conserved** (Explosion: symmetric)

Total angular momentum of homogeneous sphere:

$$J = I\omega \quad \text{where} \quad I = \frac{2}{5}MR^2$$

Neutron Stars: Rotation, II

During SN collapse, **angular momentum is conserved** (Explosion: symmetric)

Total angular momentum of homogeneous sphere:

$$J = I\omega \quad \text{where} \quad I = \frac{2}{5}MR^2$$

Angular momentum conservation ($J_{\text{before}} = J_{\text{NS}}$):

$$\frac{2}{5}M_{\text{before}}R_{\text{before}}^2\omega_{\text{before}} = \frac{2}{5}M_{\text{NS}}R_{\text{NS}}^2\omega_{\text{NS}}$$

or (assume $M_{\text{NS}} = M_{\text{before}}$):

$$\omega_{\text{NS}} = \left(\frac{R_{\text{before}}}{R_{\text{NS}}}\right)^2 \omega_{\text{before}} \quad \text{or} \quad P_{\text{NS}} = \left(\frac{R_{\text{NS}}}{R_{\text{before}}}\right)^2 P_{\text{before}}$$

(where P : rotation period)

Neutron Stars: Rotation, III

During SN collapse, **angular momentum is conserved** (Explosion: symmetric)

Total angular momentum of homogeneous sphere:

$$J = I\omega \quad \text{where} \quad I = \frac{2}{5}MR^2$$

Angular momentum conservation ($J_{\text{before}} = J_{\text{NS}}$):

$$\frac{2}{5}M_{\text{before}}R_{\text{before}}^2\omega_{\text{before}} = \frac{2}{5}M_{\text{NS}}R_{\text{NS}}^2\omega_{\text{NS}}$$

or (assume $M_{\text{NS}} = M_{\text{before}}$):

$$\omega_{\text{NS}} = \left(\frac{R_{\text{before}}}{R_{\text{NS}}}\right)^2 \omega_{\text{before}} \quad \text{or} \quad P_{\text{NS}} = \left(\frac{R_{\text{NS}}}{R_{\text{before}}}\right)^2 P_{\text{before}}$$


(where P : rotation period)

Example: $R_{\text{before}} = 700000 \text{ km}$ (sun), $R_{\text{NS}} = 15 \text{ km}$, $P_{\text{Sun}} = 27 \text{ d} \implies P_{\text{NS}} = 0.001 \text{ s}$

Neutron Stars are extremely fast rotators.

close to break-up speed!

Neutron Stars: Pulsars



Another conserved observable:

magnetic flux: $\Phi = BR^2$

magnetic field after SN:

$$B_{\text{NS}} = \left(\frac{R_{\text{before}}}{R_{\text{NS}}} \right)^2 B_{\text{before}}$$

⇒ **neutron stars have strong magnetic fields** (typical: $B \sim 10^6 \dots 10^8 \text{ T}$)

Radio pulsars are fast rotating neutron stars with strong magnetic fields.

Black Holes, I

Neutron stars also have upper mass limit: **Oppenheimer Volkoff limit**.

Detailed mass limit unknown, causality considerations give $M \sim 3 M_{\odot}$ (for “stiff equation of state” the sound speed becomes greater than speed of light at this mass)

Compact objects with mass above Oppenheimer Volkoff limit: **Black Holes**

More conservative astronomers: “Black Hole Candidates”.

Black Holes, II

Rev. John Michell: *Phil. Trans. R. Soc. London*, 74, 35–57 (1784):

VII. *On the Means of discovering the Distance, Magnitude, &c. of the Fixed Stars, in consequence of the Diminution of the Velocity of their Light, in case such a Diminution should be found to take place in any of them, and such other Data should be procured from Observations, as would be farther necessary for that Purpose. By the Rev. John Michell, B. D. F. R. S. In a Letter to Henry Cavendish, Esq. F. R. S. and A. S.*

Read November 27, 1783.

Black Holes, III

Rev. John Michell: *Phil. Trans. R. Soc. London*, **74**, 35–57 (1784):

42 *Mr. MICHELL on the Means of discovering the*

16. Hence, according to article 10, if the semi-diameter of a sphaere of the same density with the sun were to exceed that of the sun in the proportion of 500 to 1, a body falling from an infinite height towards it, would have acquired at its surface a greater velocity than that of light, and consequently, supposing light to be attracted by the same force in proportion to its vis inertiaë, with other bodies, all light emitted from such a body would be made to return towards it, by its own proper gravity.

Black Holes, IV

In more modern usage:

Total energy of a mass m :

$$E = E_{\text{pot}} + E_{\text{kin}} = -G\frac{Mm}{R} + \frac{1}{2}mv^2$$

Mass m is unbound if $E > 0$, i.e., for

$$v \geq v_{\text{escape}} = \sqrt{\frac{2GM}{R}}$$

Black Hole: Body of mass M and radius R for which $v_{\text{escape}} > c$, where c is the speed of light.

This is the case if

$$R \leq R_{\text{S}} = \frac{2GM}{c^2} \sim 3 \text{ km} \frac{M}{M_{\odot}}$$

the **Schwarzschild Radius**.

Black Holes: Accretion

Astrophysical energy sources:

1. Nuclear fusion

Reactions à la



Energy released:

Fusion produces $\sim 6 \times 10^{11} \text{ J g}^{-1}$

(i.e., $\Delta E_{\text{nuc}} \sim 0.007 m_p c^2$)

Black Holes: Accretion

Astrophysical energy sources:

1. Nuclear fusion

Reactions à la



Energy released:

Fusion produces $\sim 6 \times 10^{11} \text{ J g}^{-1}$

(i.e., $\Delta E_{\text{nuc}} \sim 0.007 m_p c^2$)

2. Gravitation

Accretion of mass m from ∞ to R_S on black hole with mass M gives

$$\Delta E_{\text{acc}} = \frac{GMm}{R_S} \text{ where } R_S = \frac{2GM}{c^2}$$

Accretion produces $\sim 10^{13} \text{ J g}^{-1}$

(i.e., $\Delta E_{\text{acc}} \sim 0.1 m_p c^2$)

Black Holes: Accretion

Astrophysical energy sources:

1. Nuclear fusion

Reactions à la



Energy released:

Fusion produces $\sim 6 \times 10^{11} \text{ J g}^{-1}$

(i.e., $\Delta E_{\text{nuc}} \sim 0.007 m_p c^2$)

2. Gravitation

Accretion of mass m from ∞ to R_S on black hole with mass M gives

$$\Delta E_{\text{acc}} = \frac{GMm}{R_S} \text{ where } R_S = \frac{2GM}{c^2}$$


Accretion produces $\sim 10^{13} \text{ J g}^{-1}$

(i.e., $\Delta E_{\text{acc}} \sim 0.1 m_p c^2$)

⇒ Accretion of material is the **most efficient** astrophysical energy source.

... thus accreting objects are the most luminous in the whole universe.

Note: energy gets radiated away from *outside* the Schwarzschild radius!



Material accretes from normal star over inner
Lagrange point, L_1 , onto compact object
 \Rightarrow Formation of an accretion disk, with temperature
 $\sim 10^7$ K
 \Rightarrow X-rays.

Summary

Stars end their lives as one of three kinds of **compact objects**:

White Dwarf: $R \sim R_{\text{Earth}}$, $\rho \sim 10^{5\text{--}6} \text{ g cm}^{-3}$

$M < 1.44 M_{\odot}$ (**Chandrasekhar Limit**)

Equilibrium between gravitation and pressure of degenerate electrons

Summary

Stars end their lives as one of three kinds of **compact objects**:

White Dwarf: $R \sim R_{\text{Earth}}$, $\rho \sim 10^{5\dots6} \text{ g cm}^{-3}$

$M < 1.44 M_{\odot}$ (**Chandrasekhar Limit**)

Equilibrium between gravitation and pressure of degenerate electrons

Neutron Star: $R \sim 10 \text{ km}$, $\rho \sim 10^{13} \dots 10^{16} \text{ g cm}^{-3}$

$1.44 M_{\odot} < M \lesssim 3 \dots 4 M_{\odot}$ (**Oppenheimer-Volkoff Limit**)

Density implies inv. β -decay ($p + e^{-} \rightarrow n$), i.e., star has high neutron content

Summary

Stars end their lives as one of three kinds of **compact objects**:

White Dwarf: $R \sim R_{\text{Earth}}$, $\rho \sim 10^{5\dots6} \text{ g cm}^{-3}$

$M < 1.44 M_{\odot}$ (**Chandrasekhar Limit**)

Equilibrium between gravitation and pressure of degenerate electrons

Neutron Star: $R \sim 10 \text{ km}$, $\rho \sim 10^{13} \dots 10^{16} \text{ g cm}^{-3}$

$1.44 M_{\odot} < M \lesssim 3 \dots 4 M_{\odot}$ (**Oppenheimer-Volkoff Limit**)

Density implies inv. β -decay ($p + e^{-} \rightarrow n$), i.e., star has high neutron content

Black Hole: Above OV-Limit no stable configuration known

\implies star collapses

\implies **Black Hole**

$M \gtrsim 4 M_{\odot}$

Event horizon at $R_S = 2GM/c^2 = 3(M/M_{\odot}) \text{ km}$ (**Schwarzschild radius**)