



The Deep Sky



© 2000, Axel Mellinger

Optical image of the whole sky

2MASS Covers the Sky




The Two Micron All Sky Survey

Infrared Processing and Analysis Center/Caltech & Univ. of Massachusetts

Infrared view of the whole sky



Milky Way, III



M83: ESO [VLT ANTU+FORS1]



Luminosity: $\sim 2 \times 10^{10} L_{\odot}$
Mass: $\sim 10^{11} M_{\odot}$ (radiating)
 $\sim 10^{12} M_{\odot}$ (total)

Stellar density:
 $\sim 0.3 M_{\odot} \text{pc}^{-3}$

$1 M_{\odot} = 2 \times 10^{30} \text{ kg}$,
 $1 L_{\odot} = 4 \times 10^{26} \text{ W}$



Milky Way, IV




Luminosity: $\sim 2 \times 10^{10} L_{\odot}$
 Mass: $\sim 10^{11} M_{\odot}$ (radiating)
 $\sim 10^{12} M_{\odot}$ (total)

Stellar density:
 $\sim 0.3 M_{\odot} \text{ pc}^{-3}$

$1 M_{\odot} = 2 \times 10^{30} \text{ kg}$,
 $1 L_{\odot} = 4 \times 10^{26} \text{ W}$

NGC 4565: W. McLaughlin




Andromeda galaxy (closest real neighbour galaxy, diam. 20kpc, distance: 675 kpc), NOAO/AURA/NSF




Virgo cluster, Burnell Schmidt telescope, NOAO/AURA/NSF

Deep looks in the universe: galaxies as building blocks



Deep image of Virgo cluster, 4 m Mayall telescope, NOAO/AURA/NSF

Edwin Hubble's Classification Scheme



SDSS


Galaxy classification via the Hubble "tuning fork diagram"



M87 (=Virgo A, note jet; E0), NOAO/AURA/NSF



M49 (E4), NOAO/AURA/NSF




M59 (E5; color image), NOAO/AURA/NSF



M86 (lenticular, S0), NOAO/AURA/NSF



Elliptical Galaxies



M60 (NGC 4649), E1, U. of Alabama

Elliptical galaxies: Classification as E_x where $x = 10(a - b)/a$ (integer part; between 0 and 7)

Ellipticals are low on dust and gas, reddish color (=old stars!), typically low luminosity and low mass ($10^6 M_{\odot}$)

Monsters: Also elliptical, from mergers in galaxy clusters (e.g., M87 in Virgo), M up to $10^{12} M_{\odot}$, designated cD.



M104 (Sa; "Sombrero galaxy), HST/NASA



Edge on Spiral NGC 4013 (Sa; NASA/HST)



M90 (Sb), NOAO/AURA/NSF



NGC 4565 (Sb, seen edge on), McLaughlin



M51 (Sc), NOAO/AURA/NSF, T. Rector



M51 (Sc; centre), HST/NASA



NGC 300 (Sc), M. Schirmer/ESO/2.2 m



Spiral Galaxies



M51 (NGC 5194 and 5195), Sc and Irr,
Kitt Peak 0.9 m

Spiral Galaxies: Elliptical nucleus plus spiral arms, designated **Sa**, **Sb**, **Sc** depending on opening angle of spiral (Sa: $\sim 10^\circ$, Sc: $\sim 20^\circ$) and dominance of nucleus.

Bluer than ellipticals.

Mass content $\sim 3 \times 10^{11} M_\odot$, with
 $M/L \sim 20$,

Gas content increases from Sa to Sc from 1% to 8%.

Spiral arms probably due to **density wave**.



M83 (SABc, ESO)



M58 (SBb), NOAO/AURA/NSF



NGC 1365 (SBb, VLT/FORS/ANTU): note old “reddish” bar, young spiral arms



Barred Galaxies



M95 (NGC 3351), SBb, INT

Barred Galaxies: Classification as **SBa**, **SBb**, **SBc** similar to S_x galaxies, but additional presence of a bar (cause of bar production and stability are still debated).

Similar masses and gas content as in normal spirals.




Large Magellanic Cloud (LMC; Irr I), Loke Kun Tan



Large Magellanic Cloud (LMC; Irr I), AURA/NOAO/NSF



Irregular Galaxies: Irr I




NGC 4449, Univ. Bonn

Irr I: no symmetry or spiral arms, bright knots of O- and B-type stars, very blue ($B - V \sim 0.5$), high dust content ($\sim 16\%$),

$M/L \sim 3$, masses vary appreciably from 10^6 to $10^{10} M_{\odot}$.


Examples: SMC, LMC
 \implies “Magellanic type irregulars”.



ESO202-G23 (VLT UT1/ISAAC/ESO)



Irregular Galaxies: Irr II



M82, HST-WFPC

Irr II: unsymmetrical and “abnormal”

⇒ All objects that do not fit in the rest of the classification: starburst galaxies, interacting galaxies, Active Galactic Nuclei, . . .



NGC 6946, T. Rector/AURA/Gemini




NGC 6946, T. Rector/AURA/Gemini

NGC 6946 is a SABc galaxy (note very small bar).



I Zwicky 18, Y. Izotov/T. Thuan/HST




I Zwicky 18, Y. Izotov/T. Thuan/HST

I Zw 18 is a irregular galaxy

(and one of the smallest galaxies known, merely 1.2 kpc across).



Hoag's Object, HST



Hoag's Object, HST

Hoag's object an irregular galaxy




NGC 1300, HST




NGC 1300, HST

NGC 1300 is a SBbc galaxy



Cen A, ESO/WFI




Cen A, ESO/WFI

Cen A is a (peculiar) S0 galaxy





AGN



NGC 3783: *linear* intensity scale




AGN

NGC 3783: *linear* intensity scale*logarithmic* intensity scale



AGN


NGC 3783: *linear* intensity scale*logarithmic* intensity scale

Active Galactic Nuclei (AGN): **supermassive black holes** ($M \sim 10^{6...8} M_{\odot}$),
accreting $1 \dots 2 M_{\odot}/\text{year}$

\Rightarrow **Luminosity** $\sim 10^{10} L_{\odot}$ (comparable to galaxy luminosity)



AGN




Structure of **active galactic nuclei** similar to galactic black holes (although somewhat scaled up. . .)

- **supermassive black hole** ($10^7 M_{\odot}$)
- **accretion disk** ($\dot{M} \sim 1 \dots 2 M_{\odot} \text{ yr}^{-1}$)
- **large luminosity** ($L \sim 10^{10} L_{\odot}$)
- **Schwarzschild radius** now $\sim 1 \text{ AU}$



AGN



Structure of **active galactic nuclei** similar to galactic black holes (although somewhat scaled up. . .)


- **supermassive black hole** ($10^7 M_{\odot}$)
- **accretion disk** ($\dot{M} \sim 1 \dots 2 M_{\odot} \text{ yr}^{-1}$)
- **large luminosity** ($L \sim 10^{10} L_{\odot}$)
- **Schwarzschild radius** now $\sim 1 \text{ AU}$
- often **relativistic jets**, where material is accelerated to the speed of light

AGN *with* jets: quasars, blazars. . .

AGN *without* jets: Seyfert galaxies



Mass Determination, I



NGC 6007 (Jansen;

<http://www.astro.rug.nl/~nfgs/>)

Spectra of galaxies: sum of all constituent spectra (mainly stars plus some contribution from nebulae).


Absorption lines show clear shift \implies **Doppler effect** due to motion of stars around centre:

$$\frac{\Delta\lambda}{\lambda} = \frac{v_r}{c} = \frac{v}{c} \sin i$$

where v_r : radial velocity, i : inclination (angle measured with respect to plane of sky).

Typical rotation speeds are a few 100 km s^{-1} .

Mass Determination, II



NGC 891 (Swaters et al., 1997, ApJ 491, 140 / Paul LeFevre, S&T Nov. 2002)




Mass Determination, III

Spiral galaxy rotation curves are flat!

“Galaxy rotation problem”, first discovered by Vera Rubin (1970)



©Astron. Soc. Pacific



← NGC 1553 (S0) (after Kormendy, 1984, ApJ 286, 116)



Rotation Curves: Interpretation



Newtonian interpretation of galaxy rotation curves:

Motion because of mass within r :

$$\frac{GM(\leq r)}{r^2} = \frac{v_{\text{rot}}^2(r)}{r}$$

such that

$$M(\leq r) = \frac{v_{\text{rot}}^2 r}{G}$$

therefore:

$v \sim \text{const.}$ implies $M(\leq r) \propto r$.

This assumption is approximately true even for nonspherical mass distributions.

NGC 891, KPNO 1.3 m
Barentine & Esquerdo



Rotation Curves: Interpretation

What mass distribution do we expect?

Intensity profile of disk in spiral galaxies can be well described by

$$I(r) = I_0 \exp(-r/h)$$

where r : distance from centre, h : “scale height”.

Luminosity emitted within radial distance r_0 :

$$L(r < r_0) = I_0 \int_0^{r_0} \exp(-r/h) 2\pi r \, dr = 2\pi I_0 \left(h^2 - \exp(-r_0/h) h(h + r_0) \right)$$


i.e., for $r_0 \rightarrow \infty$: $L(r < r_0) \rightarrow \text{const.}$

If all light comes from stars, i.e., **light traces mass**, then $M/L \sim \text{const.}$, such that $M(< r) \sim \text{const.}$ outside a certain radius and $v \propto r^{-1/2} \implies$ not what is observed!

Canonical interpretation: a large fraction of gravitating material does not emit light \implies spiral galaxies have large and massive halos made of **dark matter**, resulting in $M/L \sim 30$.



Dark Matter: MACHOS



MACHOS (Massive Compact Halo Objects): White dwarfs in the galaxy's halo

Pro:

1. very low luminosity objects
⇒ very difficult to detect
2. detected by microlensing towards SMC and LMC (see figure) ⇒ MW halo consists of 50% white dwarfs

Contra:

1. possible “self-lensing” (by stars in MW or SMC/LMC; confirmed for a few cases)
2. inferred white dwarf formation rate too high ($100 \text{ year}^{-1} \text{ Mpc}^{-3}$ instead of < 1 as previously assumed)



Dark Matter: Nonbaryonic

Nonbaryonic dark matter:

Requirements:

- **gravitating**
- **no or very weak other interaction** with baryons (=“us”)

⇒ Grab-box of elementary particle physics:

1. Neutrinos with non-zero mass

Pro: It exists, mass limits are a few eV, need only $\langle m_\nu c^2 \rangle \sim 10$ eV

Contra: ν are relativistic ($v \sim c$), this has implications for galaxy formation that make it unlikely that they form a major part of dark matter.

2. Axions ($mc^2 \sim 10^{-5...-2}$ eV) and WIMPs (weakly interacting massive particles; masses $mc^2 \sim$ GeV)

Pro: help with cosmology as well

Contra: We do not know they exist... (but they might soon be detectable)

⇒ Jury is still out, question on origin of flat rotation curves is still open.



MOND

Modified Newtonian Dynamics (Milgrom, 1983ff.; MOND): Alternative to Dark Matter

Reviews: Sanders & McGaugh, 2002, Ann. Rev. Astron. Astrophys. 40, 263; Milgrom, 2001, astro-ph/0112069

Idea: Modify Newton's Laws:

Acceleration on particle in gravitational field:

$$a = \frac{GM}{r^2} \cdot \frac{1}{\mu(a/a_0)} \quad \text{with} \quad \mu(x) \longrightarrow \begin{cases} 1 & \text{for } x \rightarrow \infty \\ x & \text{for } x \rightarrow 0 \end{cases}$$

i.e., for accelerations $a \ll a_0$, $a \longrightarrow \sqrt{GMa_0/r^2}$, giving circular motion in the limit of small accelerations:

$$\sqrt{\frac{GM(\leq r)a_0}{r^2}} = \frac{v^2}{r} \implies M(\leq r) = \frac{v^4}{Ga_0}$$

and therefore independent of r !

MOND can explain the flat rotational curves (by construction!).



MOND

Fits of rotational curves give

$$a_0 = 1.2 \times 10^{-8} \text{ cm s}^{-2}$$

and $M/L \sim 1$, so not bad!



BUT:

- where is the physics behind a_0 ?
- violation of the strong equivalence principle

(“outcome of any physical experiment is independent of where and when in the universe it is performed, and it is independent on whether the experimental apparatus is free falling or stationary”)

⇒ At the moment MOND does not seem to be a viable alternative to other theories of dark matter.

... but it shows that even today people are not afraid to attack Newton's laws, and this is good for progress of physics as a whole



after Sanders & McGaugh (2002)