



Question 1: *Lifetime of a synchrotron radiation emitting electron*

In the lecture it was shown that the radiation emitted from a relativistic electron with energy $E = \gamma m_e c^2$ is given by

$$\langle P_{\text{em}} \rangle = \frac{4}{3} \gamma^2 c \sigma_T U_B \quad (6.10)$$

A consequence of this equation is that electrons emitting synchrotron radiation loose their energy.

a) Show that the energy of a synchrotron radiation emitting electron at time t is given by

$$E(t) = \frac{E_0}{1 + t/t_{1/2}} \quad \text{with} \quad t_{1/2} = \frac{3m_e c^3}{4\sigma_T U_B} \frac{1}{E_0} \quad (\text{w1.1})$$

where $E_0 = E(t = 0)$ is the electron's initial energy.

b) Show that the life time $t_{1/2}$ can also be obtained from

$$t_{1/2} = \frac{E_0}{\langle P \rangle} \quad (\text{s1.1})$$

c) *Do this subquestion at home.* Show that for typical conditions in supernova remnants such as the Crab ($E_0 = 1 \text{ GeV}$ and $B = 10^{-4} \text{ G}$),

$$t_{1/2} = 4 \times 10^6 \text{ yr} \left(\frac{B}{10^{-4} \text{ G}} \right)^{-2} \left(\frac{E_0}{1 \text{ GeV}} \right)^{-1} \quad (\text{s1.1})$$

The Crab nebula has a size of only a few parsecs. The electrons producing the synchrotron radiation from the nebula can therefore originate from the central pulsar. The situation is different in jets from active galactic nuclei. These jets can have lengths of several 100 kpc and the typical B -fields are a few 10^{-4} G . Since the jets are significantly longer than $ct_{1/2}$, one has to infer that there is a source of energy within the jets. The physical process responsible for this source, however, is not yet known.

Hint: A good collection of cgs constants needed in this exercise can be found at <http://www.astro.wisc.edu/~dolan/constants/calc.html>.

Question 2: *Power law slope for Compton scattering*

a) A slab of thickness ℓ and electron number density n (measured in units of electrons cm^{-3}) is irradiated by light with initial intensity N_0 (where N is the number of photons per second and square-cm). Convince yourself that due to scattering, the decrease in photon number over infinitesimal distance dx is given by

$$\frac{dN}{N} = -n\sigma dx \quad (\text{w2.1})$$

where σ is the Thomson cross section. Use Eq. w2.1 to show that the number of photons emerging on the other side of the slab in the original direction of the photons is

$$N(\ell) = N_0 \exp(-\tau) \quad (\text{w2.2})$$

where $\tau = n\sigma\ell$. (*Hint:* The decrease in photon flux within an infinitely thick slab is $dN = -n\sigma N dx$.)

- b) Using Eq. (w2.2), convince yourself that the probability of a photon to travel at least an optical depth τ is

$$p(\tau) = \exp(-\tau) \quad (\text{s2.1})$$

and that the mean optical depth traveled before the photon scatters, $\langle \tau \rangle = 1$.

(*Hint*: if x has a probability density distribution $\propto p(x)$, then its mean is $\langle x \rangle = \int xp(x)dx / \int p(x)dx$.)

- c) Use the result from the previous question to show that the mean physical distance traveled in the slab, the *mean free path* l , is

$$l = \frac{1}{n\sigma} \quad (\text{s2.1})$$

- d) Show that for *small* τ the probability of a photon undergoing k scatterings before escaping the medium is approximately

$$p_k(\tau) \sim \tau^k \quad (\text{s2.1})$$

(*Hint*: First look at the probability that the photon escapes after *one* scattering and then use induction.)

- e) For Compton scattering and a seed photon energy $E_s \ll kT$, as shown in the lecture the amplification factor is

$$A \sim \frac{4kT}{mc^2} \quad (\text{s2.1})$$

Show that after k scatterings the energy of the seed photon, E_k , is approximately

$$E_k \sim E_s A^k \quad (\text{s2.2})$$

(*Hint*: proof by induction.)

- f) Using Eqs. s2.1 and s2.2, show that the emergent intensity at energy E_k is a power law

$$N(E_k) = N(E_s) \left(\frac{E_k}{E_s} \right)^{-\alpha} \quad \text{where} \quad \alpha = -\frac{\ln \tau}{\ln A} \quad (\text{s2.1})$$

(*Hint*: estimate the intensity emerging at energy E_k after k scatterings and then play with logarithms.)

Question 3: *Synchrotron self-Compton Emission and the Compton catastrophe*

Since synchrotron radiation is produced by highly energetic electrons, it is not unlikely that synchrotron photons are Compton scattered subsequent to their emission. This process is called Synchrotron self-Compton emission, and normally abbreviated SSC.

In the lectures it was shown that the power gained in Compton scattering is

$$P_{\text{compt}} = \frac{4}{3} \sigma_T c \gamma^2 \beta^2 U_{\text{rad}} \quad (\text{w3.1})$$

while the synchrotron emissivity of a plasma was

$$P_{\text{synch}} = \frac{4}{3} \sigma_T c \gamma^2 \beta^2 U_B \quad (\text{w3.2})$$

- a) The *Compton catastrophe* occurs when synchrotron photons are so violently Compton upscattered that the scattering electrons loose all of their energy, which will happen when $P_{\text{compt}} > P_{\text{synch}}$. Show that for a source of angular size θ at a distance d the Compton catastrophe will happen if the ratio

$$\frac{L_{\text{compt}}}{L_{\text{synch}}} = \frac{U_{\text{rad}}}{U_B} = \frac{8\pi F}{B^2 \theta^2 c} \quad (\text{w3.3})$$

is greater than 1. Here, F is the observed synchrotron flux from the source and $U_{\text{rad}} = L_{\text{synch}}/4\pi r^2 c$ is the energy density within the synchrotron source, which has a radius r . (*Hint*: $F = L/4\pi d^2$ and $\theta = r/d$ in the small angle approximation)

- b) The *brightness temperature*, T , is used in radio astronomy to describe the intensity, I , of a radio source. It is defined by

$$I = \frac{2\nu^2 kT}{c^2} \quad (\text{s3.1})$$

In addition, if one observes an unresolved radio source of angular size θ a flux $F = \theta^2 I$ is measured. Close to the Compton limit, the source must have a high surface brightness such that the source starts to be self-absorbed. The total flux from the source is therefore approximately $F = F(\nu_a)\nu_a$, where $F(\nu_a)$ is the source flux at the frequency ν_a , where self-absorption sets in. This frequency is approximately given by

$$\nu_a = \frac{3}{2}\nu_c \quad (\text{s3.2})$$

where $\nu_c = \omega_c/2\pi$ with $\omega_c = \gamma^2\omega_L$ being the characteristic frequency of synchrotron radiation (see lecture, Eq. 6.15). Using this information, show that

$$\frac{L_{\text{compt}}}{L_{\text{synch}}} \sim 16\pi \left(\frac{3e}{4\pi m_e^3 c^5} \right)^2 \frac{(kT)^5}{c^3} \nu_a \sim \left(\frac{T}{10^{12.1} \text{ K}} \right)^5 \frac{\nu_a}{100 \text{ GHz}} \quad (\text{s3.3})$$

Therefore, when observing opaque (=self-absorbed) sources at 100 GHz, we do not expect to see sources with brightness temperatures above 10^{12} K. This result was first noticed by Kellermann and Pauliny-Toth in 1969. If a source is found to exceed the Compton limit, this is typically due to variability and/or relativistic beaming.

Hint: Since we are observing at the frequency where self-absorption sets in, the energy of electrons radiating at that frequency is $E = kT$ where T is the observed brightness temperature.